



# The Centers

for Advanced Orthopaedics

**Lucas Wymore, MD**  
Sports Medicine  
23000 Moakley Street Suite 102  
Leonardtown MD 20650

Office Phone: 301-475-5555  
Office Fax: 301-475- 5914  
Email: lwymore@cfaortho.com

## **Rotator Cuff Repair: Postoperative Instructions**

**Diet** • Resume regular diet as soon as possible.

---

**Medication** • Take 1-2 tablets every 4-6 hours as needed for pain.

- Oxycodone    Norco    Tylenol#3
- You have been given a prescription for Phenergan. Fill this prescription **ONLY IF** you have severe nausea.
- If you had a block, begin taking pain medicine at the first sign of return of sensation. Take medicine prior to onset of pain.
- **Restart your daily medication unless you are otherwise instructed**

You will be given a prescription for pain medicine at your preoperative appointment. Take the medicine as needed according to the directions on the bottle. Possible side effects include nausea, dizziness, headache, vomiting, constipation and urinary retention. If you experience these side effects please call our office for assistance. Discontinue the medication if you develop a rash, shortness of breath, itching, or difficulty swallowing. If these symptoms become severe you should seek immediate medical attention.

**Refills on pain medication are authorized during office hours only (8am-5pm; Mon-Fri). Meds will not be refilled on weekends.**

---

**Activity**    You have been given a cooling unit to ice your shoulder. This can be used at all times for the first 1-2 weeks, and then as needed. Do not let the ice pad directly touch your skin.

Apply ice, even though bandages are thick and you may not feel the cold. Apply the ice to the shoulder 3 times per day for 20 minutes for the first 1 week until your shoulder is feeling comfortable again.

- **DO NOT** use heat.
- **DO NOT** actively (on your own) lift your operative arm away from the side of your body unless you are with your doctor, physical therapist, or athletic trainer.
- **DO NOT** lift anything with your operative hand.
- Place a pillow behind the elbow while lying down or sleeping. Sleeping in a more upright position (recliner) maybe more comfortable initially.
- Open and close your hand, flex and extend your wrist and elbow. Do each 10 times every hour that you are awake.

## SLING

- Use a sling at all times & while sleeping until your next office visit.
- You may remove your sling if you are sitting with your arm resting in your lap/on pillows

---

## Dressing Care

- Keep the dressing dry.
- You can expect some light bloody wound seepage through the bandage. **DO NOT BE ALARMED.** This is normal.

## Dressing removal:

- Remove all dressings 72 hours after the surgery. **DO NOT** remove steri-strips or sutures.
- DO NOT REMOVE** the dressings until your next office visit.

## Showering

- You may shower 3 days after surgery unless told otherwise. **DO NOT** immerse the shoulder under water and **DO NOT** rub the incision. Place new band-aids over the sutures after showering.
- Do not shower or submerge your shoulder in water. Keep the dressing clean and dry.

---

## Arthroscopic Findings

- Rotator Cuff:            Intact        Partial tear    Full tear
- Bone spur (acromion)
- Shoulder joint arthritis
- Acromioclavicular joint arthritis
- Joint lining inflammation (synovitis)
- Labrum (cartilage) tear    Anterior    Superior    Posterior
- Biceps tendon tear/inflammation
- Loose Body
- Other \_\_\_\_\_

---

## Procedure Performed

- Arthroscopic joint evaluation
- Rotator cuff debridement (trimming frayed edges)
- Rotator cuff repair through the arthroscope
- Rotator Cuff repair requiring an open incision
- Subacromial decompression (spur removal)
- Labral repair            Anterior    Superior    Posterior
- Biceps tendon tenodesis /release
- Distal Clavicle Excision (Removal of the tip of the collar bone)
- Joint debridement    Labrum    Biceps    Joint (cartilage) surface
- Other \_\_\_\_\_

---

## Physical Therapy

- You will be given a physical therapy prescription to begin in 1 to 2 days.
- You will be given a physical therapy prescription when you are seen in the office for follow-up.
- No formal physical therapy will be needed.

---

## Follow-Up

A follow-up appointment with your physician is on \_\_\_\_\_.

---

### Specific Instructions:

- Notify our office at **301-475-5555** for:
    1. A sudden increase in swelling and redness or warm at the area the surgery was performed **which is not relieved by rest, ice, and elevation.**
    2. Oral temperature greater than 101.5 degrees that is not relieved by taking 2 tylenol every 4-6 hours. **Do not exceed 3.5 grams of Tylenol over 24 hours. Note your pain medicine contains Tylenol or acetaminophen.**
    3. Excessive drainage from the incision or dressing which has not stopped 72 hours after surgery which is not relieved by applying a compressive dressing, ice and elevation
    4. Fever, chills, shortness of breath, chest pain, nausea, vomiting or other signs and symptoms that concern you.
- 

I have seen and understand these instructions.

\_\_\_\_\_  
Discharge Nurse

\_\_\_\_\_  
Patient or guardian

date \_\_\_\_\_

---