## NewportCare Medical Group

Lucas Wymore MD

40700 California Oaks Drive, Suite 205

**Sports Medicine** Office Phone:949-491-9991

Murrieta, CA 92592

Fax: 949-612-9795



## **ACL & POSTEROLATERAL CORNER RECONSTRUCTION** REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-6 weeks	Heel touch WB in brace *	<ul> <li>0-2 weeks: Locked in full extension for ambulation and sleeping</li> <li>2-6 weeks: Unlocked for ambulation 0-90, remove for sleeping**</li> </ul>	<b>0-2 weeks:</b> 0-45	Quad sets, patellar mobs, gastroc/soleus stretch
			2-6 weeks: Advance slowly 0-90	SLR w/ brace in full extension until quad strength prevents extension lag
				Side-lying hip/core
				Hamstrings avoidance until 6 wks post-op
PHASE II 6-12 weeks	Advance 25% weekly until full by 8 wks	Discontinue at 6 wks if no extension lag	Full	Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks; advance hip/core
PHASE III 12-16 weeks	Full	None	Full	Advance closed chain strengthening
				Progress proprioception activities
				Begin stairmaster, elliptical and running straight ahead at 12 weeks
PHASE IV	Full	None	Full	16 wks: Begin jumping
16-24 weeks				20 wks: Advance to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills
PHASE V > 6 months	Full	None	Full and pain- free	Gradual return to sports participation after completion
				of FSA***
				Maintenance program based on FSA

<sup>\*</sup>Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure

<sup>\*\*</sup>Brace may be removed for sleeping after first post-operative visit (day 7-10)

<sup>\*\*\*</sup>Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at 22-24 wks post-op for competitive athletes returning to play after rehab