



SICKNESS POLICY

Date: 18/04/2021

Review Date: 18/04/2022 (or when required)

Statement:

We at Pippins Preschool realise that all children have minor illnesses that do not prevent them from attending preschool however in the following circumstances children will be excluded from Pippins Preschool:

- Diarrhoea/Vomiting
- Infectious diseases

Aim:

- To reduce the spread of infection and to keep children and staff healthy and safe.

Procedure

- Please read in conjunction with our Adminstrating Medicines Policy;
- If a child arrives at preschool ill the manager will make the decision as to whether the child is fit to attend or not, if not the parent/carer will be asked to take the child home;
- If a child becomes ill during the day we will contact the parent/carer and they may be required to collect the child;
- We ask that parents/carers must inform us immediately of an infectious disease that their child has so we can alert other parents;
- By maintaining a clean environment and encouraging good hygiene we can reduce the spread of infection;
- Parents are still required to pay their fees if their child is off from preschool sick. However in the event of long term illness please speak to the manager and we will discuss alternative arrangements;
- Notifiable diseases are reported to the correct establishment (a list of notifiable diseases can be found at the bottom of this document.

CHILDHOOD DISEASES AND DISORDERS**All information below has come from the Nation Health Service (NHS) website**

Disorder	Signs and Symptoms	Incubation	Isolation from School
Bronchiolitis	Cold like to start, then irritating cough, slight high temperature, rapid wheezy breathing and difficulty eating	5-8 days <i>Very infectious</i>	Until recovered
Chickenpox and Shingles	Usually starts with a slight fever, feeling unwell and an itchy rash of flat red spots which weep and are raised, more on body than face or limbs. They scab over after 3-4 days	13 to 17 days. <i>Shingles are not infectious apart from contact with the spots</i>	Until 5 days after spots appear or when they scab over and the child feels better. Avoid contact with new-born babies, immunosuppressed persons and pregnant woman who have not had it
Conjunctivitis	Eyes are pink with discharge, watery or thick. Eyes feel gritty or sticky particularly in the morning. Spreads from one eye to both	12 hours – 3 days	Safe to come to school unless child feels unwell
Diarrhoea and Vomiting	Varies but may be abdominal pain, nausea and fever in addition to diarrhoea and vomiting	Few hours to a few days depending on cause	48 hours after last symptoms
Glandular Fever	Fatigue, sore throat, fever, faint rash ad neck gland enlargement. Possible tonsillitis or mild jaundice. Some have no signs or symptoms	4-6 weeks	Until feeling well
Hand, Foot and Mouth	Sore throat, fever and blisters on inside of the mouth and throat. These may also appear on palms, fingers and toes	3-5 days. <i>Faeces of infected person are infectious as well as discharge from nose and throat</i>	Until they are feeling well
Impetigo	Redness, particularly around the nose and mouth developing into crusty sores	4-10 days. <i>Infectious when discharge persists</i>	Once spots are healing and the child feels better
Measles	Various. Fever, conjunctivitis, cough, spots on cheeks or inside of the mouth. 3-7 days later red, blotchy rash spreading form the face and lasting for 4-7 days. Feeling very unwell	7-14 days. <i>Infectious up to 4 days after rash appears</i>	At least 4 days after the rash appears and they feel well. Avoid contact with new born babies and immunosuppressed persons
Meningitis	Severe head ache, fever, stiff neck, lowered state of consciousness, vomiting and aversion to light. Later a bluish or red blotchy rash which does go under pressure. <i>MA maybe very sudden onset and often misdiagnosed</i>	2-10 days	When fully recovered
Mumps	Fever. Swelling, tenderness to neck glands	12-25 days	5 days after symptoms first develop

Ringworm	Flat, spreading ring-shaped patches. Edges are usually reddish and crusted	4-10 days. <i>Can be caught from infected farm animals</i>	Once treatment has started
Rubella (German Measles)	Possible none in a child, fever, rash, headache and possibly joint pains and conjunctivitis	16-18 days <i>Infectious up to 4 days after rash appears</i>	5 days after the rash appears
Scarlett Fever also called Scarlatina	Symptoms of scarlet fever develop within a week of being infected. Early signs include a sore throat, a headache, a high temperature (38.3C/101F or above), swollen glands in the neck and being sick. This may be followed by a rash on the body, a red face and a white or red tongue.	Up to 7 days before symptoms start	24 hours after starting antibiotics
Slapped Cheek Disease	A high temperature, fever and nasal discharge, a bright red rash like the mark left by a slap appears on the cheeks, over the next 2 to 4 days a lacy rash spreads to the torso and limbs,	A few days before the rash appears. Children are longer contagious when the rash appears	After the rash has appeared
Whooping Cough	Cold like symptoms, followed about a week later by coughing bouts that last for a few minutes and are worse at night, may become red faced, may bring up thick mucus which can make you sick.	6 days after the cold like symptoms to 3 weeks after the coughing starts	5 days from the start of antibiotic treatment

List of notifiable diseases:

Diseases notifiable to local authority proper officers under the Health Protection (Notification) Regulations 2010:

- Acute encephalitis
- Acute infectious hepatitis
- Acute meningitis
- Acute poliomyelitis
- Anthrax
- Botulism
- Brucellosis
- Cholera
- COVID-19
- Diphtheria
- Enteric fever (typhoid or paratyphoid fever)
- Food poisoning
- Haemolytic uraemic syndrome (HUS)
- Infectious bloody diarrhoea
- Invasive group A streptococcal disease
- Legionnaires' disease
- Leprosy
- Malaria
- Measles
- Meningococcal septicaemia
- Mumps
- Plague
- Rabies
- Rubella
- Severe Acute Respiratory Syndrome (SARS)
- Scarlet fever
- Smallpox
- Tetanus
- Tuberculosis
- Typhus
- Viral haemorrhagic fever (VHF)
- Whooping cough
- Yellow fever